arXiv:math/0702739v1 [math.AG] 25 Feb 2007

Extended Hilbert's Nullstellensatz

Keqin Liu
Department of Mathematics
The University of British Columbia
Vancouver, BC
Canada, V6T 1Z2

March 17, 2006

Abstract

We prove the extended Hilbert's Nullstellensatz in the context of Hu-Liu polynomial trirings.

Which kinds of noncommutative rings are suitable for extending algebra geometry? Different attempts have been made to answer this question, but a satisfactory answer is still in hiding. My attempt at answering this question comes from the trivial extension of a ring by a bimodule over the ring. The trivial extension $R \rhd\!< S$ of a ring R by a R-bimodule ${}_R S_R$ has been used in both algebra geometry and commutative algebras for a long time ([1] and [5]). Even the R-bimodule ${}_R S_R$ is itself a ring, the multiplicative structure on the ring ${}_R S_R$ does not play any role in the trivial extension $R \rhd\!< S$, and the researchers who make use of the trivial extension $R \rhd\!< S$ have not paid attension to the multiplicative structure on the ring ${}_R S_R$. Simply speaking, my idea of choosing a class of noncommutative rings is not to forget the multiplicative structure on the ring ${}_R S_R$ while using the trivial extension $R \rhd\!< S$. If we combine the ring product on S with the bimodule actions on ${}_R S_R$ by using the Hu-Liu triassociative law, then we get a triring structure on the trivial extension $R \rhd\!< S$, which was introduced in [2]. In particular, if R is a commutative ring and the R-bimodule ${}_R S_R$ is also a commutative ring, then the resulting triring on $R \rhd\!< S$ is called a Hu-Liu triring. A Hu-Liu triring $R \rhd\!< S$ is a noncommutative ring with respect to the ring product on the trivial extension $R \rhd\!< S$ if the left R-module ${}_R S$ is different from the right R-module S_R. These kinds of Hu-Liu trirings are a class of noncommutative rings which are very close to commutative rings.

Based on the curiosity to extend algebraic geometry in the context of Hu-Liu trirings, I started on the study of affine trialgebraic sets in Chapter 5 of [2]. This paper is the continuation of the study. The main result is the extended Hilbert's Nullstellensatz.

Throughout this paper, the rings considered are assumed to have an identity, and a ring R is also denoted by $(\, R,\, +,\, \cdot\,)$ to emphasize that $\cdot$ is the ring product on the ring R.

We begin this paper by recalling some basic definitions and facts from [2]. A ring $(\, R,\, +,\, \cdot\,)$ is called a $\mathcal{Z}_2$**-ring** if there exist two subgroups R_0 and R_1 of the additive group of R, called the **even part** and **odd part** of R respectively, such that $R = R_0 \oplus R_1$ (as Abelian groups) and

$$R_0R_0 \subseteq R_0, \quad R_0R_1 + R_1R_0 \subseteq R_1, \quad R_1R_1 = 0.$$

A $\mathcal{Z}_2$-ring is nothing more than a trivial extension of a ring by a bimodule over the ring. Recently, $\mathcal{Z}_2$-rings or trivial extensions of a ring by a bimodule have been used to generalize the Lie correspondence, to extend the Frobenius' Theorem about the finite dimensional division real associative algebras, and to introduce new generalizations of Jordan algebras and Malcev algebras in [3] and [4].

Let $(\, R = R_0 \oplus R_1,\, +,\, \cdot\,)$ be a $\mathcal{Z}_2$-ring. R is called a **triring** if exists a binary operation $\sharp$ on the odd part R_1 of R such that $(\, R_1,\, +,\, \sharp\,)$ is a ring, and the two associative products $\cdot$ and $\sharp$ satisfy the **Hu-Liu triassociative law**:

$$x(\alpha \,\sharp\, \beta) = (x\alpha) \,\sharp\, \beta \quad \text{and} \quad (\alpha \,\sharp\, \beta)x = \alpha \,\sharp\, (\beta x), \tag{1}$$

where $x \in R$ and $\alpha,\, \beta \in R_1$. A triring $R = R_0 \oplus R_1$ is sometimes denoted by

$$(\, R,\, +,\, \cdot,\, \sharp\,) \quad \text{or} \quad (\, R = R_0 \oplus R_1,\, +,\, \cdot,\, \sharp\,),$$

where the associative product $\sharp$ on the odd part R_1 is called the **local product**, and the identity $1^\sharp$ of the ring $(\, R_1,\, +,\, \sharp\,)$ is called the **local identity** of the triring R. A subset I of a triring $(\, R,\, +,\, \cdot,\, \sharp\,)$ is called a **triideal** of R if I is a graded ideal of the $\mathcal{Z}_2$-ring $R = R_0 \oplus R_1$, and $I_1 = I \cap R_1$ is an ideal of the ring $(\, R_1,\, +,\, \sharp\,)$.

In order to clear up the relation between trivial extensions and Hu-Liu trirings, we need the following

Definition 1 Let R be a ring. A R-bimodule ${}_RS_R$ is called a **sharp bimodule** over R or a **sharp R-bimodule** if $(\, S,\, +,\, \sharp\,)$ is a ring and (1) holds for $x \in R$ and $\alpha,\, \beta \in S$.

Trirings are nothing more than trivial extensions of a ring by a sharp bimodule over the ring. In fact, if $(\, R = R_0 \oplus R_1,\, +,\, \cdot,\, \sharp\,)$ is a triring, then $R = R_0 \rhd\!< R_1$ is the trivial extension of the ring R_0 by the sharp R_0-bimodule ${}_{R_0}(R_1)_{R_0}$. Conversely, if $R \rhd\!< S$ is the trivial extension of a ring R by a sharp R-bimodule ${}_RS_R$, then

$$(\, R \rhd\!< S = (R \rhd\!< S)_0 \oplus (R \rhd\!< S)_1,\, +,\, \cdot,\, \sharp\,)$$

is a triring, where the local product $\sharp$ is the ring product carried by the ring $(S, +, \sharp)$, and the even part $(R \rhd\!< S)_0$ and the odd part $(R \rhd\!< S)_1$ are given by

$$(R \rhd\!< S)_0 := (R, 0) \quad \textit{and} \quad (R \rhd\!< S)_1 := (0, S).$$

We now recall the following definition from [2].

Definition 2 Let $(R = R_0 \oplus R_1, +, \cdot, \sharp)$ be a triring.

(i) R is called a **commutative triring** if both $(R, +, \cdot)$ and $(R_1, +, \sharp)$ are commutative rings.

(ii) R is called a **Hu-Liu triring** if both $(R_0, +, \cdot)$ and $(R_1, +, \sharp)$ are commutative rings.

(iii) R is called an **algebraic closed** 3**-trifield** if both $(R_0, +, \cdot)$ and $(R_1, +, \sharp)$ are algebraic closed fields, and $R_1 = R_0 1^\sharp = 1^\sharp R_0$, where $1^\sharp$ is the local identity of the triring R.

Commutative trirings are a class of commutative rings with many zero divisors. Since the zero divisors in a commutative triring are controlled nicely by the local product, the theory of commutative algebra can be extended satisfactorily in the context of commutative trirings. For example, the book [6] can be rewritten in the context of commutative trirings.

Let $(\mathbf{k} = \mathbf{k}_0 \oplus \mathbf{k}_1, +, \cdot, \sharp)$ be an algebraic closed 3-trifield with a local identity $1^\sharp$, where $\mathbf{k}_0$ and $\mathbf{k}_1$ are the even part and the odd part of $\mathbf{k}$ respectively. Let $\mathbf{k}^\sharp[n] := \mathbf{k}^\sharp[x_{(1)}, x_{(2)}, \cdots, x_{(n)}]$ be the Hu-Liu polynomial triring[1] over $\mathbf{k}$ in n indeterminates $x_{(1)}, x_{(2)}, \cdots, x_{(n)}$. The Hu-Liu polynomial triring $\mathbf{k}^\sharp[n] = \mathbf{k}^\sharp[x_{(1)}, x_{(2)}, \cdots, x_{(n)}]$ is a Hu-Liu triring, its even part

$$\mathbf{k}^\sharp[n]_0 = \mathbf{k}_0[x_{(1)0}, x_{(2)0}, \cdots, x_{(n)0}]$$

is the polynomial ring over the algebraic closed field $(\mathbf{k}_0, +, \cdot)$ in n indeterminates $x_{(1)0}, x_{(2)0}, \cdots, x_{(n)0}$, and its odd part

$$\mathbf{k}^\sharp[n]_1 = \mathbf{k}_1[x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, x_{(1)1}, \cdots, x_{(n)1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}]$$

is the polynomial ring over the algebraic closed field $(\mathbf{k}_1, +, \sharp)$ in $3n$ indeterminates $x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, x_{(1)1}, \cdots, x_{(n)1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}$. Let $\mathbf{A}^n_{\mathbf{k}}$ be the affine n-trispace over $\mathbf{k}$, where $\mathbf{A}^n_{\mathbf{k}}$ is defined by

$$\mathbf{A}^n_{\mathbf{k}} := \{ (a_{(1)}, a_{(2)}, \cdots, a_{(n)}) \mid a_{(j)} \in \mathbf{k} \text{ for } 1 \leq j \leq n \}.$$

---

[1] See Section 4.1 in [2] for the construction of the Hu-Liu polynomial triring.

If $a = (a_{(1)},\, a_{(2)},\, \cdots,\, a_{(n)}) \in \mathbf{A}^n_{\mathbf{k}}$, we define

$$\begin{aligned} a_0 : &= (a_{(1)0},\, a_{(2)0},\, \cdots,\, a_{(n)0}) \in \mathbf{A}^n_{\mathbf{k}_0}, \\ a_0 1^\sharp : &= (a_{(1)0}1^\sharp,\, a_{(2)0}1^\sharp,\, \cdots,\, a_{(n)0}1^\sharp), \\ a_1 : &= (a_{(1)1},\, a_{(2)1},\, \cdots,\, a_{(n)1}), \\ 1^\sharp a_0 : &= (1^\sharp a_{(1)0},\, 1^\sharp a_{(2)0},\, \cdots,\, 1^\sharp a_{(n)0}), \end{aligned}$$

where $a_{(j)} = a_{(j)0} + a_{(j)1}$, $a_{(j)0} \in \mathbf{k}_0$ and $a_{(j)1} \in \mathbf{k}_1$ are the even component and the odd component of $a_{(j)}$ for $1 \le j \le n$. Let $J = J_0 \oplus J_1$ be a triideal of $\mathbf{k}^\sharp[n]$. The even affine trialgebraic set $V_0^\sharp(J)$ and the odd affine trialgebraic set $V_1^\sharp(J)$ determined by the triideal J are defined by

$$V_0^\sharp(J) := \{\, a \in \mathbf{A}^n_{\mathbf{k}} \,|\, F_0(a_0) = 0 \text{ for all } F_0 \in J_0 \,\}$$

and

$$V_1^\sharp(J) := \{\, a \in \mathbf{A}^n_{\mathbf{k}} \,|\, F_1(a_0 1^\sharp,\, a_1,\, 1^\sharp a_0) = 0 \text{ for all } F_1 \in J_1 \,\}.$$

Clearly, $V_1^\sharp(J) \subseteq V_0^\sharp(J)$. Other basic properties of even and odd affine trialgebraic sets are given in Proposition 5.1 of [2].

Let V_0 and V_1 be a pair of subsets of the affine n-trispace $\mathbf{A}^n_{\mathbf{k}}$, where $V_1 \subseteq V_0$. Define $I^\sharp(V_0, V_1)$ by

$$I^\sharp(V_0, V_1) := \left\{ F_0 + F_1 \;\middle|\; \begin{array}{c} F_0 \in \mathbf{k}^\sharp[n]_0,\; F_1 \in \mathbf{k}^\sharp[n]_1, \\ F_0(a_0) = 0 \text{ for all } a \in V_0, \text{ and} \\ F_1(b_0 1^\sharp,\, b_1,\, 1^\sharp b_0) = 0 \text{ for all } b \in V_1 \end{array} \right\}.$$

Then $I^\sharp(V_0, V_1)$ is a triideal of $\mathbf{k}^\sharp[n]$.

The next proposition gives the main result of this paper.

Proposition (**The Extended Hilbert's Nullstellensatz**) Let $\mathbf{k} = \mathbf{k}_0 \oplus \mathbf{k}_1$ be an algebraic closed 3-trifield. If J is a triideal of the Hu-Liu polynomial triring $\mathbf{k}^\sharp[n] = \mathbf{k}^\sharp[x_{(1)}, x_{(2)}, \cdots, x_{(n)}]$, then $I^\sharp(V_0^\sharp(J), V_1^\sharp(J)) = \sqrt[\sharp]{J}$, where

$$\sqrt[\sharp]{J} := \left\{ G_0 + G_1 \;\middle|\; \begin{array}{c} G_0 \in \mathbf{k}^\sharp[n]_0,\; G_1 \in \mathbf{k}^\sharp[n]_1, \text{ and} \\ G_0^m \in J_0 \text{ and } G_1^{\sharp\, n} \in J_1 \text{ for some } m,\, n \in \mathcal{Z}_{>0} \end{array} \right\}$$

is the graded nilradical of J, and $G_1^{\sharp n} := \underbrace{G_1 \,\sharp\, G_1 \,\sharp\, \cdots \,\sharp\, G_1}_{n}$.

Proof Let $G_0 + G_1$ be an element of $\sqrt[\sharp]{J} = \sqrt{J_0} \oplus \sqrt{J_1}$, where $G_0 \in \sqrt{J_0}$ and $G_1 \in \sqrt{J_1}$. Then $G_0^m \in J_0$ and $G_1^{\sharp\, n} \in J_1$ for some $m,\, n \in \mathcal{Z}_{>0}$. Hence, we have

$$\Big(G_0(a_0)\Big)^m = G_0^m(a_0) = 0 \quad \text{for all } a \in V_0^\sharp(J)$$

and

$$\Big(G_1(b_0 1^\sharp,\, b_1,\, 1^\sharp b_0)\Big)^{\sharp\, n} = G_1^{\sharp\, n}(b_0 1^\sharp,\, b_1,\, 1^\sharp b_0) = 0 \quad \text{for all } b \in V_1^\sharp(J).$$

It follows that $G_0(a_0) = 0$ for all $a \in V_0^\sharp(J)$ and $G_1(b_0 1^\sharp,\, b_1,\, 1^\sharp b_0) = 0$ for all $b \in V_1^\sharp(J)$. Thus, $G_0 + G_1 \in I^\sharp(V_0^\sharp(J), V_1^\sharp(J))$. This proves

$$\sqrt[\sharp]{J} \subseteq I^\sharp(V_0^\sharp(J), V_1^\sharp(J)). \tag{2}$$

Conversely, let $F = F_0 + F_1 \in I^\sharp(V_0^\sharp(J), V_1^\sharp(J))$ with $F_i \in \mathbf{k}^\sharp[n]_i$ for $i = 0$ and 1, then

$$F_0(a_0) = 0 \quad \text{for all } a_0 \in V(J_0) \subseteq V_0^\sharp(J) \tag{3}$$

and

$$F_1(b_0 1^\sharp,\, b_1,\, 1^\sharp b_0) = 0 \quad \text{for all } b \in V_1^\sharp(J), \tag{4}$$

where $V(J_0)$ is the ordinary affine algebraic set in the affine n-space $\mathbf{A}^n_{\mathbf{k}_0}$. By (3), we have $F_0 \in I(V(J_0))$, where $I(V(J_0))$ is the ideal determined by $V(J_0)$. Using Hilbert's Nullstellensats in the polynomial ring

$$\mathbf{k}^\sharp[n]_0 = \mathbf{k}_0[x_{(1)0}, x_{(2)0}, \cdots, x_{(n)0}],$$

we get

$$F_0 \in I(V(J_0)) = \sqrt{J_0} \subseteq \sqrt[\sharp]{J}. \tag{5}$$

If $F_1 = 0$, then $F = F_0 \in \sqrt[\sharp]{J}$ by (5). Hence, we get

$$F = F_0 + F_1 \in I^\sharp(V_0^\sharp(J), V_1^\sharp(J)) \text{ and } F_1 = 0 \Longrightarrow F \in \sqrt[\sharp]{J}. \tag{6}$$

If $F_1 \neq 0$, we consider the Hu-Liu polynomial trirring

$$R := \mathbf{k}^\sharp[n+1] = \Big(\mathbf{k}^\sharp[n]\Big)^\sharp[x_{(n+1)}] = \mathbf{k}^\sharp[x_{(1)}, \cdots, x_{(n)}, x_{(n+1)}]$$

over $\mathbf{k}$ in $n+1$ indeterminates $x_{(1)}, \cdots, x_{(n)}, x_{(n+1)}$. Let

$$U := J_0 R_0 \oplus \Big(J_1 \,\sharp\, R_1 + (x_{(n+1)1} \,\sharp\, F_1 - 1^\sharp) \,\sharp\, R_1\Big),$$

where R_0 and R_1 are the even part and the odd parts of R respectively. Clearly, U is a triideal of R and $J = J_0 \oplus J_1 \subseteq U$. If

$$U_1 = J_1 \,\sharp\, R_1 + (x_{(n+1)1} \,\sharp\, F_1 - 1^\sharp) \,\sharp\, R_1 \neq R_1,$$

then $V_1^\sharp(U) \neq \emptyset$ by Proposition 5.2 (ii) in [2]. Thus, there would exist

$$c = (c_{(1)}, \cdots, c_{(n)}, c_{(n+1)}) \in \mathbf{A}^{n+1}_{\mathbf{k}}$$

such that

$$\mathbf{g}_1(c_0 1^\sharp,\, c_1,\, 1^\sharp c_0) = 0 \quad \text{for all } \mathbf{g}_1 \in U_1, \tag{7}$$

which would imply that

$$\mathbf{f}_1(d_0 1^\sharp,\, d_1,\, 1^\sharp d_0) = 0 \quad \text{for all } \mathbf{f}_1 \in J_1, \tag{8}$$

where $d = (c_{(1)}, \cdots, c_{(n)}) \in \mathbf{A}^n_{\mathbf{k}}$. It follows from (8) that

$$d = (c_{(1)}, \cdots, c_{(n)}) \in V_1^\sharp(J). \tag{9}$$

By (4), (7) and (9), we get

$$\begin{aligned} 0 &= (x_{(n+1)1} \sharp F_1 - 1^\sharp)(c) = c_{(n+1)1} \sharp F_1(c_0 1^\sharp,\, c_1,\, 1^\sharp c_0) - 1^\sharp \\ &= c_{(n+1)1} \sharp F_1(d_0 1^\sharp,\, d_1,\, 1^\sharp d_0) - 1^\sharp = c_{(n+1)1} \sharp 0 - 1^\sharp = -1^\sharp, \end{aligned}$$

which is impossible. This proves that

$$U_1 = J_1 \sharp R_1 + (x_{(n+1)1} \sharp F_1 - 1^\sharp) \sharp R_1 = R_1.$$

Therefore, the local identity $1^\sharp$ can be expressed as

$$1^\sharp = \sum_{j=1}^{m} F_{(j)} \sharp \mathbf{g}_{(j)} + (x_{(n+1)1} \sharp F_1 - 1^\sharp) \sharp \mathbf{h}, \tag{10}$$

where $m \in \mathcal{Z}_{>0}$, $F_{(j)} \in J_1$, $\mathbf{g}_{(j)} \in R_1$, $\mathbf{h} \in R_1$ and $1 \leq j \leq m$. Let

$$\mathbf{k}_1(3n) := \mathbf{k}_1\Big(x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, x_{(1)1}, \cdots, x_{(n)1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}\Big)$$

be the fractional field of the polynomial ring $\mathbf{k}_1[3n]$ over the field $(\,\mathbf{k}_1,\, +,\, \sharp\,)$ in $3n$ indeterminates $x_{(1)0}1^\sharp$, $\cdots$, $x_{(n)0}1^\sharp$, $x_{(1)1}$, $\cdots$, $x_{(n)1}$, $1^\sharp x_{(1)0}$, $\cdots$, $1^\sharp x_{(n)0}$, where

$$\mathbf{k}_1[3n] := \mathbf{k}_1\Big[x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, x_{(1)1}, \cdots, x_{(n)1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}\Big].$$

By the universal property of the polynomial ring

$$\mathbf{k}_1[3(n+1)] := \Big(\mathbf{k}_1[3n]\Big)\Big[x_{(n+1)0}1^\sharp, x_{(n+1)1}, 1^\sharp x_{(n+1)0}\Big],$$

there exists a ring homomorphism $\psi : \mathbf{k}_1[3(n+1)] \to \mathbf{k}_1(3n)$ such that

$$\psi\Big|\Big(\mathbf{k}_1[3n]\Big) = \text{ the identity map}$$

and

$$\psi\Big(x_{(n+1)0}1^\sharp\Big) = 0 = \psi\Big(1^\sharp x_{(n+1)0}\Big), \quad \psi\Big(x_{(n+1)1}\Big) = \frac{1}{F_1}.$$

After applying ψ to (10), we get

$$\begin{aligned} 1^\sharp = \psi(1^\sharp) = \sum_{j=1}^{m} \psi\left(F_{(j)}\right) \sharp \psi\Big(&\mathbf{g}_{(j)}\Big(x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, x_{(n+1)0}1^\sharp, \\ &x_{(1)1}, \cdots, x_{(n)1}, x_{(n+1)1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}, 1^\sharp x_{(n+1)0}\Big)\Big) \\ = \sum_{j=1}^{m} F_{(j)} \sharp \mathbf{g}_{(j)}\Big(&x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, 0, x_{(1)1}, \cdots, x_{(n)1}, \frac{1}{F_1}, \\ &1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}, 0\Big). \end{aligned} \tag{11}$$

Let s_j be the degree of

$$\mathbf{g}_{(j)}\Big(x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, 0, x_{(1)1}, \cdots, x_{(n)1}, \frac{1}{F_1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}, 0\Big)$$

when considered as a polynomial in $\dfrac{1}{F_1}$ with coefficients in $\mathbf{k}_1[3n]$. Then

$$(F_1)^{\sharp s} \,\sharp\, \mathbf{g}_{(j)}\Big(x_{(1)0}1^\sharp, \cdots, x_{(n)0}1^\sharp, 0, x_{(1)1}, \cdots, x_{(n)1}, \frac{1}{F_1}, 1^\sharp x_{(1)0}, \cdots, 1^\sharp x_{(n)0}, 0\Big)$$

is in $\mathbf{k}_1[3n]$, where $s := \max\{\, s_j \,|\, 1 \le j \le m \,\}$. Using this fact and multiplying both sides of (11) by $(F_1)^{\sharp s}$, we get

$$(F_1)^{\sharp s} \in \sum_{j=1}^{m} F_{(j)} \,\sharp\, (\mathbf{k}_1[3n]) \subseteq J_1. \tag{12}$$

By (5) and (12), we have

$$F = F_0 + F_1 \in I^\sharp(V_0^\sharp(J), V_1^\sharp(J)) \text{ and } F_1 \neq 0 \Longrightarrow F \in \sqrt[\sharp]{J}. \tag{13}$$

It follows from (6) and (13) that

$$\sqrt[\sharp]{J} \supseteq I^\sharp(V_0^\sharp(J), V_1^\sharp(J)). \tag{14}$$

The extended Hilbert's Nullstellensatz now follows from (2) and (14).

□

References

[1] Robert M. Fossum, Phillip A. Griffith and Idun Reiten, *Trivial Extensions of Abelian Categories*, Lecture Notes in Mathematics, Vol.456, Springer-Verlag, Berlin, 1975.

[2] Keqin Liu, *Introduction to Trirings*, Research Monographs in Mathematics **2**, 153 Publishing, 2006.

[3] Keqin Liu, *Number-like objects and the extended Lie correspondence*, arXiv: math.RA/0604179 v1 8 Apr 2006.

[4] Keqin Liu, *Generalizations of Jordan algebras and Malcev algebras*, arXiv: math.RA/0606628 v1 24 Jun 2006.

[5] Hideyuki Matsumura, *Commutative Algebra (second edition)*, The Benjamin/Cummings Publishing Company, Inc., 1980.

[6] Rodney Y. Sharp, *Steps in Commutative Algebra*, London Mathematical Society Student Texts **19**, Cambridge University Press, 1990.